

Year 2012

VCE

Mathematical Methods

CAS

Trial Examination 1



KILBAHA MULTIMEDIA PUBLISHING
PO BOX 2227
KEW VIC 3101
AUSTRALIA

TEL: (03) 9817 5374
FAX: (03) 9817 4334
kilbaha@gmail.com
<http://kilbaha.com.au>

© KILBAHA PTY LTD 2012

<http://kilbaha.com.au>

IMPORTANT COPYRIGHT NOTICE

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Pty Ltd.
 - The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
 - For authorised copying within Australia please check that your institution has a licence from Copyright Agency Limited. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.
 - Teachers and students are reminded that for the purposes of school requirements and external assessments, students must submit work that is clearly their own.
 - Schools which purchase a licence to use this material may distribute this electronic file to the students at the school for their exclusive use. This distribution can be done either on an Intranet Server or on media for the use on stand-alone computers.
 - Schools which purchase a licence to use this material may distribute this printed file to the students at the school for their exclusive use.
-
- **The Word file (if supplied) is for use ONLY within the school.**
 - **It may be modified to suit the school syllabus and for teaching purposes.**
 - **All modified versions of the file must carry this copyright notice.**
 - **Commercial use of this material is expressly prohibited.**

**Victorian Certificate of Education
2012**

STUDENT NUMBER

Figures
Words

Letter

--

MATHEMATICAL METHOD CAS

Trial Written Examination 1

Reading time: 15 minutes

Total writing time: 1 hour

QUESTION AND ANSWER BOOK

Structure of book

<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
11	11	40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers.
- Students are NOT permitted to bring into the examination room: notes of any kind, blank sheets of paper, white out liquid/tape or a calculator of any type.

Materials supplied

- Question and answer book of 14 pages with a detachable sheet of miscellaneous formulas at the end of this booklet.
- Working space is provided throughout the booklet.

Instructions

- Detach the formula sheet from the end of this book during reading time.
- Write your **student number** in the space provided above on this page.
- All written responses must be in English.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

Instructions

Answer **all** questions in the spaces provided.

In all questions where a numerical answer is required an exact value must be given unless otherwise specified.

In questions where more than one mark is available, appropriate working **must** be shown. Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Question 1

- a. Differentiate $\sqrt{16-x^3}$ with respect to x .

2 marks

- b. Hence find $\int \frac{x^2}{\sqrt{16-x^3}} dx$

1 mark

Question 2

Find the general solution of $4\cos\left(\frac{\pi x}{3}\right) - 2 = 0$.

2 marks

Question 3

The function f has the rule $f(x) = \log_e(x+3)$ and the function g has the rule $g(x) = 5 + 2x - x^2$. State the maximal domain for which $f(g(x))$ is defined.

2 marks

Question 4

The image of the line $4x + y = 3$ under the transformation $T : R^2 \rightarrow R^2$ of the plane defined by $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} 2 & 3 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} -4 \\ 2 \end{bmatrix}$, is given by $ax + by = k$, find the values of a , b and k .

3 marks

Question 5

The distance q cm, that an image is from a certain lens in terms of p cm, the distance of the object from the lens, is given by $q = \frac{10p}{p-10}$.

- i. Show that the rate of change of distance that an image is from a certain lens, with respect to the distance of the object from the lens is given by $\frac{dq}{dp} = \frac{-100}{(p-10)^2}$

1 mark

- ii. Find using a linear approximation, the change in the distance of the image from the lens when the distance of the object from the lens changes from 15 to 15.1 cm.

2 marks

- iii. If the object distance is increasing at a rate 0.25 cm/sec, how fast is the image distance changing, when the distance from the object is 15 cm.

2 marks

Question 6

The probability density function of a continuous random variable X is given by

$$f(x) = \begin{cases} kx(3-x) & \text{for } 0 \leq x \leq 3 \\ 0 & \text{elsewhere} \end{cases}$$

- i. Show that $k = \frac{2}{9}$.

1 mark

- ii. Find $\Pr(X > 2 | X > 1)$.

3 marks

Question 7

A discrete random variable X has a probability distribution given by

X	1	2
$\Pr(X = x)$	$\log_8(k+1)$	$\log_8(2k-4)$

- i. Find the possible values of k .

3 marks


- ii. Find $E(X)$, giving your answer as a fraction.

2 marks

Question 8

Part of the graph of $y = a(x-h)^3 + k$ is shown. The graph crosses the x -axis at $x = 4$, and has a stationary point at $x = 2$. The shaded area is the area bounded by the graph of $y = a(x-h)^3 + k$ and the coordinate axes, this area is equal to 64 units^2 .

Find the values of a , h and k .



4 marks

Question 10

The speed v , in metres per second, of an object moving in a straight line is given by a

function of time t , in seconds, where $v(t) = \frac{24}{\sqrt{4t+9}}$ where $t \geq 0$.

Find the distance travelled in metres by the object in the first 4 seconds.

2 marks


Question 11

Given the function $f : R \setminus \{2\} \rightarrow R$ $f(x) = \frac{3x-5}{x-2}$.

- i. Express $f(x)$ in the form $\frac{a}{x+b} + c$.

1 mark

- ii. Sketch the graph of the function $f : \mathbb{R} \setminus \{2\} \rightarrow \mathbb{R}$ $f(x) = \frac{3x-5}{x-2}$, stating the coordinates of all axial intercepts, and the equations of any asymptotes.



2 marks

- iii. Find the inverse function f^{-1} .

2 marks

END OF QUESTION AND ANSWER BOOKLET**END OF EXAMINATION****© KILBAHA PTY LTD 2012**

MATHEMATICAL METHODS CAS

Written examination 1

FORMULA SHEET

Directions to students

Detach this formula sheet during reading time.

This formula sheet is provided for your reference.

Mathematical Methods CAS Formulas

Mensuration

area of a trapezium: $\frac{1}{2}(a+b)h$

volume of a pyramid: $\frac{1}{3}Ah$

curved surface area of a cylinder: $2\pi rh$

volume of a sphere: $\frac{4}{3}\pi r^3$

volume of a cylinder: $\pi r^2 h$

area of triangle: $\frac{1}{2}bc \sin(A)$

volume of a cone: $\frac{1}{3}\pi r^2 h$

Calculus

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

$$\int x^n dx = \frac{1}{n+1}x^{n+1} + c, n \neq -1$$

$$\frac{d}{dx}(e^{ax}) = ae^{ax}$$

$$\int e^{ax} dx = \frac{1}{a}e^{ax} + c$$

$$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$$

$$\int \frac{1}{x} dx = \log_e|x| + c$$

$$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$$

$$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$$

$$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$$

$$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$$

$$\frac{d}{dx}(\tan(ax)) = \frac{a}{\cos^2(ax)} = a \sec^2(ax)$$

product rule: $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$

quotient rule: $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$

Chain rule: $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

approximation: $f(x+h) \approx f(x) + h f'(x)$

Probability

$$\Pr(A) = 1 - \Pr(A')$$

$$\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$$

$$\Pr(A/B) = \frac{\Pr(A \cap B)}{\Pr(B)}$$

Transition Matrices $S_n = T^n \times S_0$

mean: $\mu = E(X)$

variance: $\text{var}(X) = \sigma^2 = E((X - \mu)^2) = E(X^2) - \mu^2$

probability distribution		mean	variance
discrete	$\Pr(X = x) = p(x)$	$\mu = \sum x p(x)$	$\sigma^2 = \sum (x - \mu)^2 p(x)$
continuous	$\Pr(a < X < b) = \int_a^b f(x) dx$	$\mu = \int_{-\infty}^{\infty} x f(x) dx$	$\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$